

Anti-hyperlipidemia of garlic by reducing the level of total cholesterol and low-density lipoprotein

A meta-analysis

Yue-E Sun, PhD, Weidong Wang, PhD*, Jie Qin, MS

Abstract

Background: This study aimed to understand the impact of garlic on improving blood lipids using a meta-analysis.

Methods: A literature search of the PubMed, EMBASE, and Cochrane Library databases was performed using keywords such as “garlic” and “hypercholesterolemia,” and the deadline “July 14 (th), 2017.” After extracting relevant details, each selected literature was evaluated for quality according to the quality evaluation criteria of bias risk recommended by Cochrane Collaboration recommendations and heterogeneity tests were performed. Standardized mean difference (SMD) and 95% confidence interval (CI) were evaluated using R 3.12 software. The publication bias was assessed using Egger method.

Results: A total of 14 eligible papers published from 1981 to 2016 were included. The quality of the literatures was of moderate to high qualities. The values of TC (SMD = -1.26, 95% CI, -1.86 to -0.66), low-density lipoprotein (LDL) (SMD = -1.07, 95% CI, -1.67 to -0.47), and high-density lipoprotein (HDL) (SMD = 0.50, 95% CI, 0.06–0.94) after taking garlic in the experimental group and the control group have statistical significance, while there was no significant difference of TG in the 2 groups (SMD = -0.16, 95% CI, -0.87–0.55). However, the result of HDL was reversed when removed some of the literatures. No significant publication bias among the eligible studies with values of TC ($P = .0625$), LDL ($P = .0770$), HDL ($P = .2293$), and TG ($P = .3436$).

Conclusion: Garlic can reduce the level of TC and LDL instead of HDL and TG, indicating the ability of anti-hyperlipidemia.

Abbreviations: BG = aged black garlic, BMI = body mass index, CI = confidence interval, CVD = cardiovascular disease, GO = garlic oil, GP = garlic powder, HDL = high-density lipoprotein, HMG-CoA = 3-hydroxy-3-methylglutaryl-coenzyme A, LDL = low-density lipoprotein, PRISMA = Preferred Reporting Items for Systematic Reviews and Meta-Analyses, SMD = standardized mean difference, TC = total cholesterol, TG = triglyceride.

Keywords: cardiovascular disease, garlic, high-density lipoprotein, hypercholesterolemia, low-density lipoprotein, total cholesterol, triglyceride

High Lights

1. Garlic had anti-hyperlipidemia ability.
2. Intake of garlic induced the level of TC and LDL.
3. The level of TG in serum had no obvious difference with and without garlic treatment.

1. Introduction

Recently, cardiovascular disease (CVD), a complex and multi-factorial disease, remains one of the serious diseases that threaten human health worldwide with increasing incidence and mortality year by year.^[1] Unfortunately, 17 million people die to CVD each year and it is estimated to reach 24.8 million in 2030 in the world.^[2] The most important risk factors of CVD are hypertension, high cholesterol, alcohol intake, and tobacco usage, etc. according to the World Health Organization

Editor: Fadi Khasawneh.

Ethics committee: This is a meta-analysis manuscript, there is no ethics committee or institutional review board approved by the study.

This work was supported by National Natural Science Foundation of China (Program No. 31301535), Six Talent Peaks project in Jiangsu Province (Program No. NY-167), and 333 Project of Jiangsu Province (Program No. BRA2017289).

Competing Interests: The authors declare that they have no competing interests.

College of Food Engineering, Xuzhou University of Technology, Xuzhou, Jiangsu Province, China.

* Correspondence: Weidong Wang, College of Food Engineering, Xuzhou University of Technology, No.2 Lishui Road, Xincheng District, Xuzhou, Jiangsu Province, 221111, China (e-mail: wwd.123@163.com).

Copyright © 2018 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Medicine (2018) 97:18(e0255)

Received: 16 August 2017 / Received in final form: 8 February 2018 / Accepted: 1 March 2018

<http://dx.doi.org/10.1097/MD.00000000000010255>

statistics.^[3] Dyslipidemia including high total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and triglyceride (TG), as well as decreased high-density lipoprotein cholesterol (HDL-C) are important causes of CVD.^[4] It is well known that an increase in LDL-C concentration is a risk factor for CVD and is therefore considered to be the primary goal of CVD prevention and treatment.^[5] However, nowadays, most of hypolipidemic drugs have relatively great side effects during the treatment of CVD. Thus, a certain component from food with slight side-effect or free of side effects has been paid more and more attentions on the treatment of hyperlipidemia.

Garlic, a perennial herb, plays an important medicinal and dietary role throughout the history. Garlic is used in numerous forms such as extracted oil, powdered garlic tablets, or raw garlic.^[6] Ancient Indian and Chinese medicine recommend that garlic can be used to help respiration and digestion, and to treat leprosy and parasitic diseases.^[7] Besides, garlic plays important roles in the treatment of many diseases such as infectious diseases (as antibiotic), gynecologic diseases, toothache, arthritis, chronic cough.^[7] Because of the widespread effects of garlic in maintaining good health, it has attracted particular attention of modern medicine. Importantly, the favorable effects are mainly attributed to diminution of risk factors for CVDs, antimicrobial effect, antioxidant effect, reduction of cancer risk.^[8,9] The protective mechanisms of the beneficial effects of garlic in CVDs may be achieved by suppressing LDL oxidation, increasing HDL, as well as decreasing TC and TG.^[10–12] A class of sulfur-containing organic compounds (including S-allylcysteine, alliin, ajoene, diallyl disulfide) in garlic has pharmacological effects.^[13,14]

The effect of garlic in anti-hyperlipidemia has been confirmed in some animal studies, but all of them have small size of samples. Besides, a meta-analysis was performed in 2013 by Ried et al,^[15] but the studies published by Jung et al,^[5] Ahmad Alobaidi,^[16] and Negar et al^[6] are not included. Thus, in this study to increase the sample size, we used meta-analysis to understand the impact of garlic on improving blood lipids. Our study might provide a basis for clinical research.

2. Materials and methods

This meta-analysis was performed based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.^[17]

2.1. Data sources

The related clinical researches were obtained from the electronic databases PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/>), Embase (<http://www.embase.com>), and Cochrane Library (<http://www.cochranelibrary.com>) with garlic (“ALLIUM SATIVUM” OR “Garlic”) and hypercholesterolemia (“Hypercholesterolemia” OR “Hypercholesterolaemia” OR “Hypercholesterolemic” OR “Hyperlipidemic” OR “Hyperlipidaemia” OR “Hyperlipidemia” OR “Dyslipidemia” OR “Dyslipidemias”) as searched keywords; and a deadline of “July 14 (th); 2017.”

2.2. Eligibility criteria

Articles meeting the following criteria were selected (based on the PICOS principle): published English literatures to study the efficacy of garlic in the treatment of hypercholesterolemia (P); participants in the experimental group was hyperlipidemia

patients treated with garlic (I) and that in the control group was hyperlipidemia patients with placebo (C) treatment; the outcomes of the study included the initial values as well as the values after treatment of TC (serum total cholesterol), LDL (low density lipoprotein), HDL (high density lipoprotein), and TG (triglyceride), and the amount of changes was included (O); the study type was randomized parallel study (study design: parallel; crossover study: excluded) (S). The following articles were removed: studies with incomplete data or can't be used for statistical analysis; literatures such as reviews, reports, comments, and letters. Besides, if multiple literatures were repeatedly published or multiple literatures studied based on the same population data, only the latest research or the research with complete information was included.

2.3. Data extraction and quality evaluation of literature

Data were independently extracted from the included literatures by 2 reviewers and included details such as: first author, published year, area of study, year of study, the type and dose of garlic, follow-up time, number as well as general demographic data (e.g., sex ratio, age composition, body mass index [BMI], etc.) of inclusion in the garlic group and the control group; the initial values as well as the values after treatment of TC, LDL, HDL, and TG in the garlic group and the control group.

The aggregate quality of the included studies was evaluated according to the quality evaluation criteria of bias risk, recommended by Cochrane Collaboration recommendations.^[18] A third reviewer should join in and discuss with the other 2 reviewers to get an agreement if there was disagreement in the process of data extraction and quality evaluation.

2.4. Statistical analysis

The meta-analysis of direct comparison was conducted using R 3.12 software (R Foundation for Statistical Computing, Beijing1, China, “meta” package), SMD (standardized mean difference), and 95% confidence interval (CI) were used to show the effect index of quantitative data. The heterogeneity test between studies was assessed based on the Q test^[19] and the I^2 statistic, and $P < .05$ or $I^2 > 50\%$ was used as the heterogeneity threshold. The random effect model was chosen when they had significant heterogeneity; otherwise, the fixed-effect model was chosen to pool the data.^[20] The publication bias was assessed using Egger method. Finally, the sensitivity analysis was performed by examining the effect of this document on the overall SMD value by ignoring a document each time.

3. Results

3.1. Characteristics of the selected literature

A total of 956 articles (243 articles came from PubMed database, 645 articles came from Embase database, 68 ones came from Cochrane Library) were identified based on the literature search criteria. Among them, 207 articles were repeated, and 618 articles were irrelevant after reading title and abstract. In addition, 117 articles (including 15 letters, 11 case series/report, 28 reviews, 31 animal study, 13 descriptive studies, 12 non-RCT, and 7 reduplicative studies) of the remaining 131 articles were removed by reviewing full text. Finally, 14 eligible papers which were published from 1981 to 2016 were included (Fig. 1).^[5,6,16,21–31]




Figure 1. Literature search and study selection.

Totally, 1093 hyperlipidemia patients were included in the study, including 527 cases in the experimental group and 566 cases in the control group. Most of the areas of these studies were India, Canada, the United States, and South Korea; the types of garlic were aged black garlic (BG), garlic oil (GO), and garlic powder (GP) with the doses ranged from 0.3 to 20g/d; the patients were followed up for 4 weeks to 10 months; the baseline characteristics between the experimental group and the control group were comparable in terms of age, sex, and BMI/Body

weight (Tables 1 and 2). The results of RCT quality evaluation showed that the quality of the literatures we incorporated were of moderate to high qualities (Fig. 2A, B).

3.2. Combination of quantitative data of meta-analysis

The values of TC, LDL, HDL, and TG in the experimental group and the control group after taking garlic were analyzed by meta-analysis. Based on the heterogeneity test results ($P < .05$, $I^2 > 50\%$), all results including TC ($P < .0001$, $I^2 = 94\%$), LDL ($P < .0001$, $I^2 = 94.2\%$), HDL ($P < .0001$, $I^2 = 91.1\%$), and TG ($P < .0001$, $I^2 = 95.9\%$) were pooled using random model.

The results of meta-analysis showed that the values of TC, LDL, and HDL after taking garlic in the experimental group and the control group have statistical significance. Among that, garlic treatment decreased the values of TC (SMD = -1.26, 95% CI, -1.86 to -0.66, Fig. 3) and LDL (SMD = -1.07, 95% CI, -1.67 to -0.47, Fig. 4) while increased that of HDL (SMD = 0.50, 95% CI, 0.06-0.94, Fig. 5), indicating the important role of garlic in changing blood lipids. However, there was no significant difference of TG in the 2 groups (SMD = -0.16, 95% CI, -0.87-0.55, Fig. 6).

3.3. Publication bias

Egger test was used to assess the publication biases in the analyses of TC, LDL, HDL, and TG values. Our results showed no significant publication bias among the eligible studies with values of TC ($P = .0625$), LDL ($P = .0770$), HDL ($P = .2293$), and TG ($P = .3436$, Table 3).

Table 1

Characteristics of the included literatures.

Author	Year	Study location	Type	Dose	Duration	Group	N	Sex (M/F)	Age	BMI/Body weight
Adler AJ	1997	Canada	GP	900mg/d	12 w	Garlic	12	12 (M)	45.9±12.6	27.2±3.0
						Placebo	11	11 (M)	45.4±9.8	26.4±3.5
Ahmad Alobaidi AH	2014	Iraq	GO	500mg/d	4 w	Garlic	150	66/84	46.13±5.25	26.38±5.91
						Placebo	150	61/89	45.03±7.52	27.15±6.73
Ashraf R	2005	Pakistan	GP	300mg/d	12 w	Garlic	35	15/20	60±5.04	62.2±10.45kg
						Placebo	35	17/18	58±5.80	63±9.80kg
Aslani N	2016	Iran	Garlic	20g/d	8 w	Garlic	27	14/13	45.3±9.3	28.1±5.9
						Placebo	28	16/12	39.3±6.2	27.2±3.2
Bordia A	1981	India	GO	15mg/d	10 m	Garlic	20	NA	54.7 (44-62)	NA
						Placebo	62		53.0 (42-60)	NA
Gardner CDa	2007	USA	Others	4g/d	6 m	Raw garlic	49	22/27	49±9	25±3
						Placebo	48	24/24	49±9	25±3
Jain AK	1993	USA	GP	900mg/d	12 w	Garlic	20	11/9	48±15	78±17kg
						Placebo	22	8/14	55±9	77±14kg
Jung ES	2014	Korea	ABG	6g/d	12 w	Garlic	30	12/18	50.13±9.21	64.69±9.88kg
						Placebo	30	8/22	50.83±8.04	63.57±9.76kg
Kannar D	2001	Australia	GP	880mg/d	12 w	Garlic	22	12/10	52.6±10.4	24.5±2.6
						Placebo	24	13/11	57.4±9.0	28.4±6.3
Peleg A	2003	Israel	GP	22.4mg/d	16 w	Garlic	18	8/10	52.4±7.5	69.3±11.8kg
						Placebo	21	9/12	54.7±7.5	70.1±8.4kg
Satitvipawee P	2003	Thailand	Others	333mg/d	12 w	Garlic	70	23/47	47±6.6	24.6±3.3
						Placebo	66	25/41	47±6.0	24.3±3.4
Sobenin LA	2008	Russia	GP	600mg/d	12 w	Garlic	23	42 (M)	51.7±2.0	26.6±0.5
						Placebo	19		51.7±2.0	27.0±0.7
Sobenin LA	2010	Russia	GP	300mg/d	12 m	Garlic	26	14/12	56.7±1.8	27.0±0.9
						Placebo	25	14/11	56.3±1.7	27.9±0.9
Superko HR	2000	USA	GP	900mg/d	12 w	Garlic	25	NA	53±107	163±30 lbs
						Placebo	25			

Garlic type: a=raw garlic group, ABG=aged black garlic, BMI=body mass index, GO=garlic oil, GP=garlic powder, M/F=Male/Female.

Table 2**Detailed data of the included literatures.**

Author	Year	Group	N	TC		LDL		HDL		Triglyceride	
				Base, mg/dL	Posttreatment, mg/dL	Base, mg/dL	Posttreatment, mg/dL	Base, mg/dL	Posttreatment, mg/dL	Base, mg/dL	Posttreatment, mg/dL
Adler AJ	1997	Garlic	12	6.54±0.25a	5.79±0.23a	4.39±0.18a	3.77±0.24a	1.26±0.06a	1.29±0.08a	1.98±0.32a	1.85±0.37a
		Placebo	11	6.46±0.26a	6.49±0.31a	4.32±0.25a	4.26±0.31a	1.20±0.10a	1.26±0.11a	1.95±0.49a	1.94±0.51a
AlhmadAlobaidi AH	2014	Garlic	150	217±46.1	176±34.7	139.4±28.5	98.3±30.13	37.8±13.9	45.9±15.1	199±39.7	159±33.1
		Placebo	150	213.6±41.5	198.6±35.4	134.7±33.6	119.1±35.4	38.1±13.5	42.1±13.7	201±46.6	187±37.6
Ashraf R	2005	Garlic	35	228.23±4.54	200.77±5.07	163.57±4.66	133.42±4.61	38±1.73	41.35±1.31	202.03±8.59	203.45±8.81
		Placebo	33	220.45±2.25	218.34±3.05	167±3.37	164.3±3.56	36.58±3.45	37.2±3.86	200.6±5.05	198.58±5.25
Aslani N	2016	Garlic	27	234.4±26.5	215.3±23	119.9±26.8	105.1±23	37.2±7.4	41±8.6	195.9±123.7	162±98.6
		Placebo	28	239±18.8	243.1±14.2	119.2±19.0	127.2±22.1	41.3±6.5	41.3±7.8	135.6±64.9	151.1±67.1
Bordia A	1981	Garlic	20	298.4±22.7	228.0±13.5	NA	NA	17.0±1.1	30.0±2.6	NA	NA
		Placebo	62	280.1±25.1	282.4±18.0	NA	NA	17.8±1.3	17.4±1.8	NA	NA
Gardner Cda	2007	Raw garlic	49	226±18	NA	151±15	142±22	58±15	58±14	97±44	95±49
		Placebo	48	228±21	NA	150±14	133±21	54±14	52±13	122±54	134±74
Jain AK	1993	Garlic	20	262±35	247±40	188±37	168±43	47±12	46±13	151±81	165±86
		Placebo	22	276±34	274±29	191±34	185±25	49±14	50±17	195±112	199±101
Jung ES	2014	Garlic	30	241.07±23.97	233.50±24.63	150.64±14.12	155.75±21.94	46.86±9.40	50.36±8.85	139.79±54.55	120.21±63.61
		Placebo	30	228.93±20.01	227.33±32.51	150.11±15.65	156.33±29.83	50.81±9.19	50.48±9.76	124.11±47.65	128.11±65.50
Kannar D	2001	Garlic	22	7.5±0.8a	7.4±1.1a	5.3±0.9a	5.4±1.1a	1.34±0.34a	1.33±0.39a	2.0±1.3a	1.8±1.0a
		Placebo	24	7.6±0.9a	7.1±0.9a	5.3±0.9a	4.9±0.9a	1.35±0.49a	1.26±0.47a	2.2±1.6a	2.3±2.4a
Peleg A	2003	Garlic	18	262.6±25.3	259.6±38.6	172.7±18.8	171.0±28.3	54.0±11.9	49.8±13.3	179.5±69.6	231.4±13.9
		Placebo	21	275.4±23.9	267.7±29.6	186.6±16.8	182.0±23.5	54.9±15.8	54.0±11.9	169.6±70.7	157.9±44.3
Saithipawee P	2003	Garlic	70	6.65±0.89a	6.59±0.93a	4.52±0.86a	4.52±0.77a	1.50±0.37a	1.45±0.28a	1.19±0.02a	1.17±0.02a
		Placebo	66	6.85±0.83a	6.80±0.90a	4.63±0.90a	4.65±0.83a	1.55±0.26a	1.47±0.26a	1.27±0.02a	1.26±0.02a
Sobenin LA	2008	Garlic	23	6.97±0.20a	6.41±0.22a	5.00±0.17a	4.37±0.20a	1.06±0.07a	1.17±0.09a	2.00±0.26a	1.91±0.21a
		Placebo	19	7.04±0.18a	7.24±0.18a	4.93±0.18a	5.07±0.16a	1.20±0.09a	1.16±0.10a	2.25±0.20a	2.06±0.22a
Sobenin LA	2010	Garlic	26	269.2±11.5	235.7±8.4	185.9±9.3	155.6±7.8	50.9±3.6	51.5±3.1	162.1±19.5	142.9±16.5
		Placebo	25	252.5±9.1	242.0±6.9	178.3±9.1	169.9±7.2	48.7±2.5	50.7±2.2	127.8±9.7	106.9±6.8
Superko HR	2000	Garlic	25	250±29	248±23	169±25	167±25	51.3±11.5	51.2±10.1	145±54	141±57.2
		Placebo	25	239±23	248±22	162±18	159±19	51.9±12.8	52.1±11.0	128±63	150±111.5

a=mmol/L, HDL=high density lipoprotein, LDL=low density lipoprotein, TC=serum total cholesterol, TG=triglyceride.




Figure 2. Quality assessments of the included studies. A. Sensitivity and specificity of the included studies. B. Bias risk of the included studies. "+": low risk of bias; "-": high risk of bias, and "?": unclear risk of bias.

3.4. Sensitivity analysis

The sensitivity analysis results revealed that any of the literature can't change the results of TC, LDL, and TG, indicating that the results of TC, LDL, and TG were stable. However, the result of HDL was reversed when removed some of the literatures (Ashraf^[23] and Bordia^[24]).

4. Discussion

In order to evaluate the reliability of previous studies, a meta-analysis evaluating the hypolipidemic effect of garlic was conducted in this study. Our finding revealed that the values of TC, LDL, and HDL after taking garlic in the experimental group and the control group have statistical significance. However, there was no significant difference of TG in the 2 groups.

Garlic is gained substantial interest by many researchers because of its impact on lipid levels.^[32,33] Garlic is discovered has multiple useful cardiovascular effects including reduction in cholesterol and TG, lowering of blood pressure, and enhancement of fibrinolytic activity.^[34] Many studies have demonstrated that different extracts of garlic can alone lower the level of serum TC, LDL, and TG in humans and rodents.^[35,36] Similarly, Maha et al^[37] have revealed that the level of plasma TC and LDL-C can be decreased by adding 8% raw garlic into the diet of rats. In 1993, Warshafsky et al^[38] have proved that intake of garlic can reduce the cholesterol level by about 10%. Besides, combination of lemon juice and garlic obviously decreased serum TC, LDL-C, and blood pressure.^[6] Nevertheless, someone have indicated that garlic powder doesn't reduce cholesterol levels, which may due to the loss of active compound(s) during processing.^[39] Garlic may decrease the absorption of cholesterol, and the synthesis of cholesterol and fatty acid, and thereby reduces the level of cholesterol.^[40] The human enzymes required in cholesterol biosynthesis such as squalene monooxygenase and HMG-CoA (3-hydroxy-3-methylglutaryl-coenzyme A) reductase can be inhibited by garlic and various constituents.^[41-45] Garlic may decrease the level of LDL-C by reduction of hepatic cholesterol 7 α -hydroxylase, HMG-CoA reductase, pentose-phosphate pathway activities,^[46] enhancement of bile acid excretion, microsomal triglyceride transfer protein,^[47] cholesteryl ester transfer protein activity,^[48] bile acid excretion,^[38] and prohibiting hepatic fatty acid synthesis,^[49] which was conducted by allicin and/or other components in garlic.^[41] Our finding showed the significant differences of TC, LDL, and HDL between the experimental group and the control group, nevertheless, the level of LG was not obviously different. Our results might suggest that the ability of garlic to lower cholesterol and LDL was better than that to lower TG.

Based on the heterogeneity test results, all results including TC, LDL, HDL, and TG were pooled using random model indicating the exist of significant heterogeneity. The appearance of heterogeneity may be due to the following reasons: garlic types were different between different articles, the patients in studies published by Ahmad Alobaidi^[16] and Bordia^[24] were treated via garlic oil, and that in the study of Jung et al^[5] were given via aged black garlic; the dose and duration were dissimilar between different articles, indicating the dose and duration can affect the result of anti-hyperlipidemia; altered detected method might be the influencing factor; the unit of BMI/body weight was dissimilar, Ashraf,^[23] Jain et al,^[21] Jung et al,^[5] and Peleg et al^[28] evaluated the physical quality via body weight, while others evaluated that by BMI, thus, it is difficult to compare the physical quality, suggesting physical quality might also affect the results. Besides, in the present study, when removed the literatures published by Ashraf^[23] and Bordia,^[24] the result of HDL was reversed. The reason might be that garlic increased the level of HDL, and the participants with type 2 diabetes and coronary heart disease were enrolled in the two studies.




Figure 3. Comparison of TC value after garlic treatment between the experimental group and the control group. TC=serum total cholesterol.




Figure 4. Comparison of LDL value after garlic treatment between the experimental group and the control group. LDL=low density lipoprotein.




Figure 5. Comparison of HDL value after garlic treatment between the experimental group and the control group. HDL=high density lipoprotein.




Figure 6. Comparison of TG value after garlic treatment between the experimental group and the control group. TG=triglyceride.

Table 3

Results of meta-analysis.

Variable	Sample size			Test of association				Test of heterogeneity*†			Egger's test‡	
	K	Cases	Control	SMD (95% CI)	Z	P	Model	Q	P	I ² (%)	t	P
TC	13	478	516	-1.2575 [-1.8582; -0.6568]	-4.1031	<.0001	Random	198.43	<0.01	94.0	2.0722	.0625
LDL	13	507	502	-1.0705 [-1.6669; -0.4741]	-3.5178	.0004	Random	208.50	<0.01	94.2	1.9511	.0770
HDL	14	527	564	0.4986 [0.0579; 0.9394]	2.2172	.0266	Random	146.16	<0.01	91.1	1.2666	.2293
TG	13	507	502	-0.1616 [-0.8708; 0.5477]	-0.4465	.6553	Random	295.84	<0.01	95.9	0.9898	.3436

P<.05 is considered statistically significant; OR=odds ratio; CI=confidence interval; K=number of studies combined.
 *Random-effects model was used when the P for heterogeneity test <.05, otherwise the fixed-effect model was used.
 †P<.05 is considered statistically significant for Q statistics.
 ‡Egger's test to evaluate publication bias.

However, this research had some limitations as following: this study did not adjust for covariates, and no further conduct subgroup analysis due to the incomplete data of some studies; the reason why specific heterogeneity exit was not determined; the results of HDL value were unstable due to the reverse finding after removed some of the articles.

In conclusion, this study using a meta-analysis demonstrated that garlic can reduce the level of TC and LDL instead of HDL and TG, indicating the ability of anti-hyperlipidemia. Thus, it is recommended that people with high blood lipids can eat more garlic. But this result requires more high-quality studies or a larger sample size of updated meta-analysis to verify.

Author contributions

Conceptualization: Weidong Wang, Yue-E Sun.
 Data curation: Jie Qin.
 Formal analysis: Jie Qin.
 Writing – original draft: Yue-E Sun.
 Writing – review and editing: Weidong Wang.

References

- Meves H, Nagy K. Cardiovascular disease in Europe 2014: epidemiological update. Eur Heart J 2014;35:2950–9.
- Estruch R, Ros E, Salas-Salvado J, et al. Primary prevention of cardiovascular disease with a Mediterranean diet. N Engl J Med 2013; 368:1279–90.
- Wood D, Joint European Societies Task F. Established and emerging cardiovascular risk factors. Am Heart J 2001;141(2 Suppl):S49–57.
- Miller M. Niacin as a component of combination therapy for dyslipidemia. Mayo Clin Proc 2003;78:735–42.
- Jung ES, Park SH, Choi EK, et al. Reduction of blood lipid parameters by a 12-wk supplementation of aged black garlic: a randomized controlled trial. Nutrition 2014;30:1034–9.
- Negar A, Hasan EM, Gholamreza A, et al. Effect of garlic and lemon juice mixture on lipid profile and some cardiovascular risk factors in people 30-60 years old with moderate hyperlipidaemia: a randomized clinical trial. Int J Prev Med 2016;7:95.
- Bayan L, Koulivand PH, Gorji A. Garlic: a review of potential therapeutic effects. Avicenna J Phytomed 2014;4:1–4.
- Colin AL. The antioxidant mechanisms underlying the aged garlic extract- and S-allylcysteine-induced protection. Oxid Med Cell Long 2012;2012:907162.
- Aviello G, Abenavoli L, Borrelli F, et al. Garlic: empiricism or science? Nat Prod Commun 2009;4:1785–96.
- Katsuki T, Hirata K, Ishikawa H, et al. Significance of garlic and its constituents in cancer and cardiovascular disease. J Am Acad Dermatol 2006;36:S1–2.
- Gardner CD, Chatterjee LM, Carlson JJ. The effect of a garlic preparation on plasma lipid levels in moderately hypercholesterolemic adults. Atherosclerosis 2001;154:213–20.
- Ziaei S, Hantoshzadeh S, Rezasoltani P, et al. The effect of garlic tablet on plasma lipids and platelet aggregation in nulliparous pregnant at high risk of preeclampsia. Eur J Obstet Gynecol Reprod Biol 2001;99:201–6.
- Iciek M, Kwiecień I, Wlodek L. Biological properties of garlic and garlic-derived organosulfur compounds. Environ Mol Mutagen 2010;50: 247–65.
- Elbayoumy K, Sinha R, Pinto JT, et al. Cancer chemoprevention by garlic and garlic-containing sulfur and selenium compounds. J Nutr 2006;136 (3 Suppl):864S–9S.
- Ried K, Toben C, Fakler P. Effect of garlic on serum lipids: an updated meta-analysis. Nutrition reviews 2013;71:282–99.
- Ahmad Alobaidi AH. Effect of Nigella sativa and Allium sativum coadministered with simvastatin in dyslipidemia patients: a prospective,

- randomized, double-blind trial. *Antiinflamm Antiallergy Agents Med Chem* 2014;13:68–74.
- [17] Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 2009;6:e1000100.
- [18] Higgins JP, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. UK: Wiley Online Library, John Wiley & Sons, Ltd; 2008.
- [19] Lau J, Ioannidis JP, Schmid CH. Quantitative synthesis in systematic reviews. *Ann Intern Med* 1997;127:820–6.
- [20] Feng R-N, Zhao C, Sun C-H, et al. Meta-analysis of TNF 308 G/A polymorphism and type 2 diabetes mellitus. *PLoS One* 2011;6:e18480.
- [21] Jain AK, Vargas R, Gotzkowsky S, et al. Can garlic reduce levels of serum lipids? A controlled clinical study. *Am J Med* 1993;94:632–5.
- [22] Adler AJ, Holub BJ. Effect of garlic and fish-oil supplementation on serum lipid and lipoprotein concentrations in hypercholesterolemic men. *Am J Clin Nutr* 1997;65:445–50.
- [23] Ashraf R, Aamir K, Shaikh AR, et al. Effects of garlic on dyslipidemia in patients with type 2 diabetes mellitus. *J Ayub Med Coll Abbottabad* 2005;17:60–4.
- [24] Bordia A. Effect of garlic on blood lipids in patients with coronary heart disease. *Am J Clin Nutr* 1981;34:2100–3.
- [25] Gardner CD, Lawson LD, Block E, et al. Effect of raw garlic vs commercial garlic supplements on plasma lipid concentrations in adults with moderate hypercholesterolemia: a randomized clinical trial. *Arch Intern Med* 2007;167:346–53.
- [26] Sobenin A, Priyanishnikov VV, Kunnova LM, et al. The effects of time-released garlic powder tablets on multifunctional cardiovascular risk in patients with coronary artery disease. *Lipids Health Dis* 2010;9:119.
- [27] Kannar D, Wattanapenpaiboon N, Savige GS, et al. Hypocholesterolemic effect of an enteric-coated garlic supplement. *J Am Coll Nutr* 2001;20:225–31.
- [28] Peleg A, Hershovici T, Lipa R, et al. Effect of garlic on lipid profile and psychopathologic parameters in people with mild to moderate hypercholesterolemia. *Isr Med Assoc J* 2003;5:637–40.
- [29] Satitvipawee P, Rawdaree P, Indrabhakti S, et al. No effect of garlic extract supplement on serum lipid levels in hypercholesterolemic subjects. *J Med Assoc Thai* 2003;86:750–7.
- [30] Sobenin IA, Andrianova IV, Demidova ON, et al. Lipid-lowering effects of time-released garlic powder tablets in double-blinded placebo-controlled randomized study. *J Atheroscler Thromb* 2008;15:334–8.
- [31] Superko HR, Krauss RM. Garlic powder, effect on plasma lipids, postprandial lipemia, low-density lipoprotein particle size, high-density lipoprotein subclass distribution and lipoprotein(a). *J Am Coll Cardiol* 2000;35:321–6.
- [32] Bhalla K, Hwang BJ, Dewi RE, et al. Metformin prevents liver tumorigenesis by inhibiting pathways driving hepatic lipogenesis. *Cancer Prev Res (Phila)* 2012;5:544–52.
- [33] Augusti KT. Therapeutic values of onion (*Allium cepa* L.) and garlic (*Allium sativum* L). *Indian J Exp Biol* 1996;34:634–40.
- [34] Batsis JA, Lopez-Jimenez F. Cardiovascular risk assessment - From individual risk prediction to estimation of global risk and change in risk in the population. *BMC Med* 2010;8:29.
- [35] Budoff MJ, Ahmadi N, Gul KM, et al. Aged garlic extract supplemented with B vitamins, folic acid and L-arginine retards the progression of subclinical atherosclerosis: a randomized clinical trial. *Prev Med* 2009;49:101–7.
- [36] Rahman K. Garlic and aging: new insights into an old remedy. *Ageing Res Rev* 2003;2:39–56.
- [37] Elmahdi B, Maha MK, Afaf IA. The effect of fresh crushed garlic bulbs (*Allium sativum*) on plasma lipids in hypercholesterolemic rats. *Res J Anim Vet Sci* 2008;3:15–9.
- [38] Warshafsky S, Kamer RS, Sivak SL. Effect of garlic on total serum cholesterol. A meta-analysis. *Ann Intern Med* 1993;119((7 Pt 1)):599–605.
- [39] Isaacsohn JL, Moser M, Stein EA, et al. Garlic powder and plasma lipids and lipoproteins: a multicenter, randomized, placebo-controlled trial. *Arch Intern Med* 1998;158:1189–94.
- [40] Matsuura H. Saponins in garlic as modifiers of the risk of cardiovascular disease. *J Nutr* 2001;131:1000S–5S.
- [41] Gebhardt R. Multiple inhibitory effects of garlic extracts on cholesterol biosynthesis in hepatocytes. *Lipids* 1993;28:613–9.
- [42] Gupta N, Porter TD. Garlic and garlic-derived compounds inhibit human squalene monooxygenase. *J Nutr* 2001;131:1662–7.
- [43] Liu L, Yeh YY. Water-soluble organosulfur compounds of garlic inhibit fatty acid and triglyceride syntheses in cultured rat hepatocytes. *Lipids* 2001;36:395–400.
- [44] Yeh YY, Liu L. Cholesterol lowering effect of garlic extracts and organosulfur compounds: human and animal studies. *J Nutr* 2001;131:989S–93S.
- [45] Augusti KT, Chackery J, Jacob J, et al. Beneficial effects of a polar fraction of garlic (*Allium sativum* Linn) oil in rats fed with two different high fat diets. *Indian J Exp Biol* 2005;43:76–83.
- [46] Qureshi AA, Din ZZ, Abuirmeileh N, et al. Suppression of avian hepatic lipid metabolism by solvent extracts of garlic: impact on serum lipids. *J Nutr* 1983;113:1746–55.
- [47] Lin MC, Wang EJ, Lee C, et al. Garlic inhibits microsomal triglyceride transfer protein gene expression in human liver and intestinal cell lines and in rat intestine. *J Nutr* 2002;132:1165–8.
- [48] Song YS, Choi MS, Park SJ, et al. Cholesteryl ester transfer protein activity and atherogenic parameters in rabbits supplemented with cholesterol and garlic powder. *Life Sci* 2003;72:2953–64.
- [49] Chang ML, Johnson MA. Effect of garlic on carbohydrate metabolism and lipid synthesis in rats. *J Nutr* 1980;110:931–6.